

SURFACE TRIANGULATION – THE METRIC APPROACH

EMIL SAUCAN

ABSTRACT. We embark in a program of studying the problem of better approximating surfaces by triangulations (triangular meshes) by considering the approximating triangulations as finite metric spaces and the target smooth surface as their Hausdorff-Gromov limit. This allows us to define in a more natural way the relevant elements, constants and invariants s.a. principal directions and principal values, Gaussian and Mean curvature, etc. By a "natural way" we mean an intrinsic, discrete, metric definitions as opposed to approximating or paraphrasing the differentiable notions. In this way we hope to circumvent computational errors and, indeed, conceptual ones, that are often inherent to the classical, "numerical" approach. In this first study we consider the problem of determining the Gaussian curvature of a polyhedral surface, by using the *embedding curvature* in the sense of Wald (and Menger). We present two modalities of employing these definitions for the computation of Gaussian curvature.

1. INTRODUCTION

The paramount importance of triangulations of surfaces and their ubiquity in various implementations (s.a. in numerous algorithms applied in robot (and computer) vision, computer graphics and geometric modelling, with a wide range of applications from industrial ones, to biomedical engineering to cartography and astrophysics – to number just a few) has hardly to be underlined here. In consequence, determining the intrinsic proprieties of the surfaces under study, and especially computing their Gaussian curvature is essential. However Gaussian curvature is a notion that is defined for smooth surfaces only, and usually attacked with differential tools, tools that – however ingenious and learned – can hardly represent good approximations for curvature of PL -surfaces, since they are usually just discretizations of formulas developed in the smooth (i.e. of class at least C^2) case.¹

Moreover, since considering triangulations, one is faced with finite graphs, or, in many cases (when given just the vertices of the triangulation) only with finite –thus discrete – metric spaces. Therefore, the following natural questions arise: (A) Is one fully justified in employing discrete metric spaces when evaluating numerical invariants of continuous surfaces? and (B) Can one find discrete, metric equivalents of the differentiable notions, notions that are intrinsically more apt to describe the properties of the finite spaces under investigations? One is further motivated to ask the questions above, since the metric method we propose to employ have already

Date: 24.01.2004.

We would like to thank Prof. Gershon Elber of The Computer Science Department, The Technion, Haifa, who motivated and sustained this project.

¹ However one can find very scientifically sound discrete versions of Surface Curvature can be found, for instance, in [Ba2], [BCM], [C-SM].

successfully been used in the such diverse fields as Geometric Group Theory, Geometric Topology and Hyperbolic Manifolds, and Geometric Measure Theory. Their relevance to Computer Graphics in particular and Applied Mathematics in general is made even more poignant by the study of *Clouds of Points* (see [LWZL], [MD]) and also in applications in Chemistry (see [T]).

We show that the answer to both this questions is affirmative, and we focus our investigations mainly on the study of metric equivalents of the Gauss curvature. Their role is not restricted to that of being yet another discrete version of Gaussian Curvature, but permits us to attach a meaningful notion of curvature to points where the surface fails to be smooth, such as *cone points* and *critical lines*. Thus we can employ curvature in reconstructin not only smooth surface, but also surfaces with "folds", "ridges" and "facets".

This exposition is organized as follows: in Section 2 we concentrate our efforts on the theoretical level and study the Lipschitz and Gromov-Hausdorff distances between metric spaces, and show that approximating smooth surfaces by nets and triangulations is not only permissible, but is, in a way, the natural thing to do, in particular we show that every compact surface is the Gromov-Hausdorff limit of a sequence of finite graphs.² In Section 3 we introduce the best candidate for a metric (discrete) version of the classical Gauss curvature of smooth surfaces, that is the Embedding, or Wald curvature. We study its proprieties and investigate the relationship between the Wald and the Gauss curvatures, and show that for smooth surfaces they coincide, so that the Wald curvature represents a legitimate discrete candidate for approximating the Gaussian curvature of triangulated surfaces. Section 4 is dedicated to developing formulas that allow the computation of Wald curvature: first the precise ones, based upon the Cayley-Menger determinants, and then we develop (after Robinson) elementary formulas that approximate well the Embedding curvature. We conclude with three Appendices. In the first Appendix we present three metric analogues for the curvature of curves, namely the Menger, Alt and Haantjes curvatures and study their mutual relationship. Furthermore we show how to relate to these notions as metric analogues of sectional curvature and how to employ them in the evaluation of Gauss curvature of triangulated surfaces. Next we present yet another metric analogue of surfaces curvature, based, in this case, upon a the modern triangle comparison method, namely the Rinow curvature. We investigate its proprieties and show (following Kirk ([K])) that in the case under investigation the Rinow and Wald curvatures coincide (and therefore Rinow curvature also identifies to the Gauss curvature). The third and last Appendix is dedicated to the development of determinant formula for the radius of the circumscribed sphere around a tetrahedron, with a view towards applications.

2. THE HAUSDORFF-GROMOV LIMITS

2.1. Lipschitz Distance. This definition is based upon a very simple³ idea: it measures the *relative* difference between metrics, more precisely it evaluates their ratio; i.e.:

The metric spaces (X, d_X) , (Y, d_Y) are *close* iff $\exists f : X \xrightarrow{\sim} Y$ s.t. $\frac{d_Y(fx, fy)}{d_X(x, y)} \approx 1$,

² For the relevance of these notions in the study of classical curvatures convergence, see [CMS], [F].

³ That is to say: very intuitive, i.e. based upon physical measurements.

$\forall x, y \in X$.⁴

Technically, we give the following:

Definition 2.1. The map $f : (X, d_X) \rightarrow (Y, d_Y)$ is *bi-Lipschitz* iff $\exists c, C > 0$ s.t.:

$$c \cdot d_X(x, y) \leq d_Y(fx, fy) \leq C \cdot d_X(x, y).$$

Remark 2.2. The same definition applies for two different metrics d_1, d_2 on the same space X .

Definition 2.3. Given a Lipschitz map $f : X \rightarrow Y$, we define the *dilatation* of f by:

$$\text{dil } f = \sup_{x \neq y \in X} \frac{d_Y(fx, fy)}{d_X(x, y)}.$$

Remark 2.4. The dilatation represents the *minimal* Lipschitz constant of maps between X and Y .

Remark 2.5. If f is not Lipschitz, then $\text{dil } f \triangleq \infty$.

Remark 2.6. (1) f Lipschitz $\Rightarrow f$ continuous.

(2) f bi-Lipschitz $\Rightarrow f$ homeo. on its image.

Remark 2.7. We have the following results:

Proposition 2.8. Let $f, g : X \rightarrow Y$ be Lipschitz maps. Then:

(a) $g \circ f$ is Lipschitz

and

(b) $\text{dil}(g \circ f) \leq \text{dil } f \cdot \text{dil } g$

Proposition 2.9. The set $\{f : (X, d) \rightarrow \mathbb{R} \mid f \text{ Lipschitz}\}$ is a vector space.

Now we can return to our main interest and define the following notion:

Definition 2.10. Let $(X, d_X), (Y, d_Y)$ be metric spaces. Then the *Lipschitz distance* between (X, d_X) and (Y, d_Y) is defined as:

$$d_L(X, Y) = \inf_{\substack{f: X \xrightarrow{\sim} Y \\ f \text{ bi-Lip.}}} \log \max(\text{dil } f, \text{dil } f^{-1})$$

Remark 2.11. If $\neq f$ bi-Lipschitz between X and Y , then – remembering Remark 2.2 – we put $d_L(X, Y) \triangleq \infty$ (i.e. d_L is **not** suited for pairs of spaces that are **not** bi-Lipschitz equivalent.)

having defined the distance between two metric spaces we now can define the *convergence* in this metric in the following natural way:

Definition 2.12. The sequence of metric spaces $\{(X_n, d_n)\}$ convergence to the metric space $\{(X, d)\}$ iff

$$\lim_n d_L(X_n, X) = 0$$

(In this case we write: $(X_n, d_n) \xrightarrow{L} 0$).

Example 2.13. Let S_t be a family of regular surfaces, $S_t = f_t(U)$; where U is an open set, $U = \text{int } U \subseteq \mathbb{R}$; such that the family $\{f_t\}$ of parametrizations is smooth (i.e. $F : U \times \mathbb{R} \rightarrow \mathbb{R}^3 \in \mathcal{C}^1$; where $F((u, v), t) \stackrel{\text{not}}{=} f_t(u, v)$). Then $d_n(S_t, S_0) \xrightarrow{t \rightarrow 0} 0$.

⁴ Here and in the sequel "fx" etc. ... stands as a short-hand version of "f(x)".

Remark 2.14. If F is not smooth (only continuous) then we do not necessarily have that $S_{t \rightarrow 0} S_0$.

We have the following significant theorem:

Theorem 2.15. *The d_L satisfies the following conditions:*

- (a) $d_L \geq 0$;
 - (b) d_L is symmetric;
 - (c) d_L satisfies the triangle inequality;
- Moreover, if X, Y are compact, then:
- (d) $d_L(X, Y) = 0 \Leftrightarrow X \cong Y$ (i.e. X is isometric to Y);
- that is

d_L is a metric on the space of isometry classes of compact metric spaces

Remark 2.16. Let us recall the following

Definition 2.17. $(X_n, d_n) \xrightarrow{u} (X, d) \triangleq d_n \xrightarrow{u} d$ as a real function; i.e.

$$\sup_{x, y \in X} |d_n(x, y) - d(x, y)| \xrightarrow{u} 0$$

(where "u" denotes uniform convergence.)

Then $X_n \xrightarrow{u} X \Rightarrow X_n \xrightarrow{L} X$ but $X_n \xrightarrow{L} X \not\Rightarrow X_n \xrightarrow{u} X$. However, for *finite* spaces indeed $X_n \xrightarrow{u} X \Leftrightarrow X_n \xrightarrow{L} X$.

2.2. Gromov-Hausdorff distance. This is also a distance between compact metric spaces ((distinguished) up to isometry!). However it gives a *weaker* topology (In particular: it is *always finite* (even for pairs of non-homeomorphic spaces.))⁵ We start by first introducing the classical

2.2.1. Hausdorff distance.

Definition 2.18. Let $A, B \subseteq (X, d)$. We define the *Hausdorff distance* between A and B as:

$$d_H(A, B) = \inf\{r > 0 \mid A \subset U_r(B), B \subset U_r(A)\}$$

(see Fig. 1); where $U_r(A)$ is the r -neighborhood of A , $U_r(A) \triangleq \bigcup_{a \in A} B_r(a)$; (here, as usual: $B_r(a) = \{x \in X \mid d(a, x) < r\}$.)

Another (equivalent) way of defining the Hausdorff distance is as follows:

$$d_H(A, B) = \max\left\{\sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A)\right\}.$$

(see Fig. 2)

We have the following

Proposition 2.19. *Let (X, d) be a metric space. Then:*

- (a) d_H is a semi-metric (on 2^X). (i.e. $A = B \Rightarrow d_H(A, B) = 0$.)
- (b) $d_H(A, \bar{A}) = 0, \forall A \subseteq X$.
- (c) $(A = \bar{A} \text{ and } B = \bar{B}) \Rightarrow (d_H(A, B) = 0 \Leftrightarrow A = B)$.

i.e. d_H is a metric on the set of closed subsets of X .

⁵The relationship between the Lipschitz and the Hausdorff distances is akin to that between the C^0 and C^1 norms in Functional Spaces.




FIGURE 1.

Notation We put: $\mathcal{M}(X) = (\{A \subseteq X \mid A = \bar{A}\}, d_H) = 2^X/d_H$.

Remark 2.20. (1) if X is compact and if $\{A_n\}_{n \geq 1} \subseteq X$ is a sequence of compact subsets of X , then:

(a) $A_{n+1} \subseteq A_n \Rightarrow A_n \xrightarrow{H} \bigcap_{n \geq 1} A_n$.

(b) $A_n \subseteq A_{n+1} \Rightarrow A_n \xrightarrow{H} \bigcup_{n \geq 1} A_n$.

(2) For general subsets $A_n \xrightarrow{H} A \in \mathcal{M}(X)$, and

(a) $A = \{\lim_n a_n \mid a_n \in A_n; n \geq 1\}$.

(b) $A = \xrightarrow{H} \bigcap_{n \geq 1} (\overline{\bigcup_{m=n}^{\infty} A_m})$.

(3) If $A_n \xrightarrow{H} A$, and if the sets A_n are all convex, then A is convex sets.

We have the following two important results, which we present without their respective (lengthy) proofs:

Proposition 2.21. X complete $\Rightarrow \mathcal{M}(X)$ complete.

Theorem 2.22. (Blaschke) X compact $\Rightarrow \mathcal{M}(X)$ compact.




FIGURE 2.

2.3. The Gromov-Hausdorff Distance. We are now able to define the Gromov-Hausdorff distance using the following basic guide-lines: we want to get the *maximum* distance that satisfies the following two conditions:

(a) $d_{GH}(A, B) \leq d_H(A, B), \forall A, B \subset X$ (i.e. set that are close as subsets of X will still be close as abstract metric spaces);

and

(b) X isometric to $Y \iff d_{GH}(X, Y) = 0$.

Definition 2.23. Let X, Y be metric spaces. Then the *Gromov-Hausdorff distance* between X and Y is defined by:

$$d_{GH}(X, Y) \triangleq \inf d_H^Z(f(X), g(Y))$$

where the infimum is taken over all the isometric embeddings $f : X \hookrightarrow Z, g : Y \hookrightarrow Z$ into some metric space Z . (See Fig. 3).




FIGURE 3.

Remark 2.24. If $X = \mathbb{S}^2$, with the spherical metric, and $Z = \mathbb{R}^3$, with the Euclidian metric, then $f(X) \neq X$ (!)

Example 2.25. Let Y be an ε -net⁶ in X . Then $d_{GH}(X, Y) \leq \varepsilon$.

Proof Take $Z = X = X', Y = Y'$. □

Remark 2.26. It is sufficient to consider embeddings f into the disjoint union of the spaces X and Y , $X \coprod Y$.

Remark 2.27. (1) X, Y bounded $\implies d_{GH}(X, Y) \leq \infty$.

(2) If $\text{diam} X, \text{diam} Y < \infty$, then $d_{GH}(X, Y) \geq \frac{1}{2}|\text{diam} X - \text{diam} Y|$ ⁷.

⁶ **Definition** Let (X, d) be a metric space, and let $A \subset X$. A is called an ε -net iff $d(x, A) \leq \varepsilon, \forall x \in X$.

⁷ $\text{diam} X \triangleq \sup_{x, y \in X} d(x, y)$

However, the straightforward definition of d_{GH} may be difficult to implement. Therefore we would like to estimate (compute) d_{GH} by comparing distances in X vs. distances in Y (as done in the cases of uniform and Lipschitz metrics). We start by defining a *correspondence* between metric spaces: $X \longleftrightarrow Y$, given by correspondences $x \leftrightarrow y$ between points $x \in X, y \in Y$.

Remark 2.28. A correspondence is **not** necessarily a function, that is to a single x may correspond to several y -s.

We shall prove that

$$(*) \quad d_{GH}(X, Y) < r \iff \exists \text{ a correspondence } X \longleftrightarrow Y \\ \text{s.t. } (x \leftrightarrow y, x' \leftrightarrow y') \implies |d_X(x, x') - d_Y(y, y')| < 2r$$

Formally, we have:

Definition 2.29. Let X, Y denote sets. A *correspondence* $X \longleftrightarrow Y$ is a subset of the Cartesian product of X and Y : $\mathcal{R} \subset X \times Y$ s.t.

- (i) $\forall x \in X, \exists y \in Y, \text{s.t. } (x, y) \in \mathcal{R}$;
- and
- (ii) $\forall y \in Y, \exists x \in X, \text{s.t. } (x, y) \in \mathcal{R}$.

Example 2.30. Any surjective function $f : X \rightarrow Y$ represents correspondence $\mathcal{R} = \{(x, f(x))\}$.

Remark 2.31. $\mathcal{R}$ is a correspondence $\iff \exists Z$ and $\exists f : Z \rightarrow X, \exists g : Z \rightarrow Y$; f, g surjective, s.t. $\mathcal{R} = \{(f(z), g(z)) \mid z \in Z\}$.

Definition 2.32. Let $\mathcal{R}$ be a correspondence between X and Y , where X, Y are metric spaces. We Define the *distortion* of $\mathcal{R}$ by:

$$\text{dis } \mathcal{R} = \sup \{ |d_X(x, x') - d_Y(y, y')| \mid (x, y), (x', y') \in \mathcal{R} \}.$$

(See (*) .)

Remark 2.33. (1) If $\mathcal{R} = \{(x, f(x))\}$ is a correspondence induced by a surjective function $f : X \rightarrow Y$, then $\text{dis } \mathcal{R} = \text{dis } f$, where:

$$\text{dis } f \triangleq \sup_{a, b \in X} |d_Y(fa, fb) - d_X(a, b)|^8.$$

- (2) If $\mathcal{R} = \{(f(z), g(z))\}$, where $f : X \rightarrow Z, g : Y \rightarrow Z$ are surjective functions, then:

$$\text{dis } \mathcal{R} = \sup_{z, z' \in Z} \{ |d_X(fz, fz') - d_Y(gz, gz')| \}.$$

- (3) $\mathcal{R} = 0$ iff $\mathcal{R}$ is associated to an isometry.

We bring, without proof, the following theorem:

Theorem 2.34. Let X, Y be metric spaces. Then:

$$d_{GH}(X, Y) = \frac{1}{2} \inf_{\mathcal{R}} (\text{dis } \mathcal{R});$$

where the infimum is taken over all the correspondences $X \xleftrightarrow{\mathcal{R}} Y$.

⁸ Remember that any correspondence can be expressed in this functional manner.

Before bringing the next result (which is very important in determining the topology) we first introduce one more notion:

Definition 2.35. $f : X \rightarrow Y$ is called an ε -isometry ($\varepsilon > 0$), iff

- (i) $\text{dis } f \leq \varepsilon$,
- and
- (ii) $f(X)$ is an ε -net in Y .

Remark 2.36. f ε -isometry $\not\Rightarrow f$ continuous.

Corollary 2.37. Let X, Y be metric spaces and let $\varepsilon > 0$. Then:

- (i) $d_{GH}(X, Y) < \varepsilon \implies \exists 2\varepsilon$ -isometry $f : X \rightarrow Y$.
- (ii) $\exists \varepsilon$ -isometry $f : X \rightarrow Y \implies d_{GH}(X, Y) < 2\varepsilon$.

Proof (i) Let $X \xleftarrow{\mathcal{R}} Y$ s.t. $\text{dis } \mathcal{R} < 2\varepsilon$.

For any $x \in X$ and $f(x) \in Y$, choose $y = f(x)$ s.t. $(x, f(x)) \in \mathcal{R}$. Then $x \mapsto f(x)$ defines a map $f : X \rightarrow Y$. Moreover: $\text{dil } f \leq \text{dil } \mathcal{R} < \varepsilon$.

We shall prove that $f(X)$ is a 2ε -net in Y .

Indeed, let $x \in X$ and $y \in Y$ s.t. $(x, y) \in \mathcal{R}$. Then $d(y, f(x)) \leq d(x, x) + \text{dis } \mathcal{R} < 2r$, thence $d(y, f(X)) < 2r$. □

Let f be an 2ε -isometry. Define $\mathcal{R} \subset X \times Y$, $\mathcal{R} = \{(x, y) \mid d(y, f(x)) \leq \varepsilon\}$.

Then, since $f(X)$ is an ε -net it follows that $\mathcal{R}$ is a correspondence.

Then $\forall (x, y), (x', y') \in \mathcal{R}$ we have:

$$\begin{aligned} |d_Y(y, y') - d_X(x, x')| &\leq |d(fx, f'x') - d(x, x')| + d(y, fx) + d(y', f'x') \leq \text{dis } f + \varepsilon + \varepsilon \leq 3\varepsilon. \\ \implies \text{dis } \mathcal{R} &\leq 3\varepsilon \implies d_{GH}(x, y) \leq 3r/2 < 2r. \end{aligned}$$

□

The next result is of great importance (in particular so in our context):

Theorem 2.38. d_{GH} is a (finite) metric on the set of isometry classes of compact metric spaces.

Proof It suffices to prove that $d_{GH}(X, Y) = 0 \implies X \cong Y$.⁹

Indeed, let X, Y be compact spaces s.t. $d_{GH} = 0$. Then it follows from the previous Corollary (for $\varepsilon = 1/n$) that $\exists (f_n)_{n \geq 1}$, $f_n : X \rightarrow Y$ s.t. $\text{dis } f_n \xrightarrow{n} 0$.

let $S \subset X$, $\bar{S}$, $|S| = \aleph_0$. Using a Cantor-diagonal argument one easily shows that $\exists (f_{n_k})_{k \geq 1} \subset (f_n)_{n \geq 1}$ s.t. $(f_{n_k})_{k \geq 1}$ converges in $Y, \forall x \in S$. Without restricting the generality we may assume that this happens for $(f_n)_{n \geq 1}$ itself. Thus we can define a function $f : X \rightarrow Y$ by putting: $f(x) = \lim_n f_n(x)$.

But $|d(f_n x, f_n y) - d(x, y)| \leq \text{dis } f_n \xrightarrow{n} 0 \implies d(fx, fy) = \lim d(f_n x, f_n y)$. In other words $f|_S$ is an isometry. But $S = \bar{S}$, therefore this isometry can be extended to an isometry $\tilde{f}$ from X to Y . In a analogous manner one shows the existence of an isometry $\tilde{f} : X \rightarrow Y$. □

Remark 2.39. $X_n \xrightarrow{L} X \implies X_n \xrightarrow{u} X \implies X_n \xrightarrow{GH} X$.

In fact, the following relationship exists between " $\xrightarrow{L}$ " and " $\xrightarrow{GH}$ ":

Theorem 2.40. $X_n \xrightarrow{GH} X \iff \varepsilon$ -nets in $X_n \xrightarrow{L} \varepsilon$ -nets in X .

⁹We shall write: $X \cong Y$ if X is isometric to Y .

One can formulate this assertion in a more formal manner and it directly (see [G+], pg. 73). However we shall proceed in more "delicate" manner, starting with:

Definition 2.41. Let X, Y be compact metric spaces, and let $\varepsilon, \delta > 0$. X, Y are called ε - δ -approximations (of each-other) iff: $\exists \{x_i\}_{i=1}^N \subset X, \exists \{y_i\}_{i=1}^N \subset Y$ s.t.

- (i) $\{x_i\}_{i=1}^N$ is an ε -net in X and $\{y_i\}_{i=1}^N$ is an ε -net in Y ;
- (ii) $|d_X(x_i, x_j) - d_Y(y_i, y_j)| < \delta \forall i, j \in \{1, \dots, N\}$.

An $(\varepsilon, \varepsilon)$ -approximation is called, for short: an ε -approximation.

The relationship between this last definition and the Gromov-Hausdorff distance is first revealed in

Proposition 2.42. *Let X, Y be compact metric spaces. Then:*

- (1) *If Y is a (ε, δ) -approximation of X , then $d_{GH}(X, Y) < 2\varepsilon + \delta$.*
- (2) *$d_{GH}(X, Y) < \varepsilon \implies Y$ is a 5ε -approximation of X .*

Proof (1) Condition (ii) of Def. 2.41. is equivalent to $\text{dis } \mathcal{R}_{X_0 Y_0} < \delta$, where $X_0 = \{x_i\}_{i=1}^N, \{y_i\}_{i=1}^N$. But $\text{dis } \mathcal{R}_{X_0 Y_0} < \delta \implies d_{GH}(X_0, Y_0) < \delta/2$. Now, since X_0 and Y_0 are ε -nets in X , resp. Y , it follows that $d_{GH}(X, X_0) \leq \varepsilon, d_{GH}(Y, Y_0) < \varepsilon$. From here and from the $d_{GH}(X_0, Y_0) < \delta/2$ follows, by means of the triangle inequality, that $d_{GH}(X, Y) < 2\varepsilon + \delta$. $\square$

(2) By Cor. 2.37., there exists a 2ε -isometrie $f : X \rightarrow Y$. Let $X_0 = \{x_i\}_{i=1}^N$ be an ε -net, and let $y_i = f(x_i)$.

Then $|d(x_i, x_j) - d(y_i, y_j)| < 2\varepsilon < 5\varepsilon$. Therefore suffice to prove that $Y_0 = \{y_i\}_{i=1}^N$ is a 5ε -net in Y .

Indeed, if $y \in Y$, then, since $f(X)$ is an 2ε -net in Y , $\exists x \in X$ s.t. $d(y, f(x)) < 2\varepsilon$. Now, since X_0 is an ε -net in X , $\exists x_i \in X_0$, s.t. $d(x, x_i) \leq \varepsilon$.

$$\begin{aligned} \text{Therefore: } d(y, y_i) &= d(y, f(x_i)) \leq d(y, f(x)) + d(f(x), f(x_i)) \\ &\leq 2\varepsilon + d(x, x_i) + \text{dis } f \\ &\leq 2\varepsilon + \varepsilon + 2\varepsilon \leq 5\varepsilon. \end{aligned}$$

$\square$

Remark 2.43. Prop. 2.42. $\iff (X_n \xrightarrow{GH} X) \iff (\forall \varepsilon > 0, X_n \text{ is an } \varepsilon\text{-approximation, } \forall n \text{ large enough.})$

More precisely we have the following Proposition:

Proposition 2.44. *Let $X, \{X_n\}_{n=1}^\infty$ compact metric spaces. Then:*

$X_n \xrightarrow{GH} X \iff \forall \varepsilon > 0, \exists$ a finite ε -net $S \subset X$ and $\exists$ a finite ε -net $S_n \subset X_n$, s.t. $S_n \xrightarrow{GH} S$ and, moreover, $|S_n| = |S|$, for large enough n .

Proof ($\implies$) If S, S_n exist, then $\implies X_n$ is an ε -approximation of $X \xrightarrow{2.42} X_n \xrightarrow{GH} X \forall n$. $\square$

($\implies$) Let S be a finite $\varepsilon/2$ -net in X .

We construct in X_n corresponding nets S_n (to be more precise, we define: $S_n = f_n(X)$, where f_n is an ε_n -approximation, $f_n : X \rightarrow X_n, \varepsilon_n \rightarrow 0$.) Then $S_n \xrightarrow{GH} S$ and, in addition, S_n is an ε -net in S (for n large enough). $\square$

We make the following extremely important Remark:

Remark 2.45. Let $\mathcal{M}(n, k, D)$ be an n -dimensional manifold, of (sectional, Ricci) curvature $\leq k$, and s.t. $\text{diam } \mathcal{M} \geq D$. Then $(\mathcal{M}, d_{GH})$ is compact. However, it should be noted that this result doesn't hold for curvature $< k$. (only for $\text{Vol}(\mathcal{M}) \leq V_0$) and injectivity radius $\geq r_0$.

Note With the notations of the precedent Proposition, the distances in S_n converge to the distances in S , as $X_n \xrightarrow{GH} X$, therefore **The Geometric Proprieties of S_n will converge to those of S .** Thus we can use the Gromov-Hausdorff each and every time **The Geometric Proprieties of X_n can be expressed in term of a finite number of points, and, by passing to the limit, automatically obtain proprieties of X .**

A typical example is that of the *intrinsic metric* i.e. *the metric induced by a length structure (i.e. path length) by a metric on a subset (of a given metric space).* (See Fig. 4 for the classical example of surfaces in $\mathbb{R}^3$.)




FIGURE 4.

On a more formal note, we have the following characterization of intrinsic metrics:

Theorem 2.46. *Let (X, d) be a complete metric space.*

- (1) *If $\forall x, y \in X$, $\exists \frac{1}{2}xy$, then d is strictly intrinsic.*
- (2) *If $\forall x, y \in X$ and $\forall \varepsilon > 0$, $\exists$ the ε -middle of xy , then d is intrinsic.*

Where we used the following definitions and notations:

Definition 2.47. (1) Given x, y points in (X, d) , the *middle* (or *midpoint*) of the *segment* xy (more correctly: 'a midpoint between "x" and "y"') is defined as:

$$\frac{1}{2}xy = z, \quad d(x, z) = d(z, y).$$

- (2) d is called *strictly intrinsic* iff the length structure is associated with is complete.
- (3) Let d be an intrinsic metric. z is an ε -middle (or an ε -midpoint) for xy iff: $|2d(x, z) - d(x, y)| \leq \varepsilon$ and $|2d(y, z) - d(x, y)| \leq \varepsilon$.

Remark 2.48. The converse of Thm. 2.46. holds in any metric space, more precisely we have:

Proposition 2.49. *If d is an intrinsic metric, then $\frac{1}{2}xy$ exists, $\forall x, y$.*

The following Theorem shows that length spaces are closed in the GH -topology:

Theorem 2.50. *Let $\{X_n\}$ be length spaces and let X be a complete metric space s.t. $X_n \xrightarrow{GH} X$. Then X is a length space.*

Proof We have already presented the idea of the proof: it is sufficient to show that for every x, y there exist an ε -midpoint ($\forall \varepsilon > 0$).

Indeed, let n be such that $d_{GH} < \frac{\varepsilon}{10}$. Then, from the a preceding result, it follows that there exist a correspondence $X_n \xleftrightarrow{\mathcal{R}} X$ s.t. $\text{dis } \mathcal{R} < \frac{\varepsilon}{5}$.

Let $\bar{x}, \bar{y} \in X_n$, $x \xleftrightarrow{\mathcal{R}} \bar{x}$, $y \xleftrightarrow{\mathcal{R}} \bar{y}$. Since X_n is a length space, $\implies \exists \bar{z} \in X_n$ s.t. $\bar{z} = \frac{\varepsilon}{5}$ -midpoint of $\bar{x}\bar{y}$. Consider $z \in X$, $z \xleftrightarrow{\mathcal{R}} \bar{z}$. Then:

$$||xz| - \frac{1}{2}|xy|| \leq ||\bar{x}\bar{z}| - \frac{1}{2}|\bar{x}\bar{y}|| + 2\text{dis } \mathcal{R} < \frac{\varepsilon}{5} + \frac{2\varepsilon}{5} < \varepsilon.$$

(Here we write $|xy|$ instead of $d(x, y)$, etc.)

In a similar manner we show that: $||yz| - \frac{1}{2}|xy|| < \varepsilon$; i.e. ε -midpoint of xy . □

The next Theorem and its Corollary are of paramount importance:

Theorem 2.51. *Any compact length space is the GH -limit of a sequence of finite graphs.*

Proof Let ε, δ ($\delta \ll \varepsilon$) small enough, and let S be a δ -net in X .

Let $G = (V, E)$ be the graph with $V = S$ and $E = \{(x, y) \mid d(x, y) < \varepsilon\}$. we shall prove that G is an ε -approximation of X , for δ small enough (i.e. for $\delta < \frac{\varepsilon^2}{4} \text{diam}(X)$). (See Fig. 5.)




FIGURE 5.

But, since S is an ε -net both in X and in G , and since $d_G(x, y) \geq d_X(x, y)$, it is sufficient to prove that:

$$d_G(x, y) \leq d_X(x, y) + \varepsilon.$$

Let γ be the shortest path between x and y , and let $x_1, \dots, x_n \in \gamma$ s.t. $n \leq \text{length}(\gamma)/\varepsilon$ (and $|x_i, x_{i+1}| \leq \varepsilon/2$). Since $\forall x_i \exists y_i \in S$ s.t. $|x_i, y_i| \leq \delta$, it follows that $|y_i y_{i+1}| \leq |x_i x_{i+1}| + 2\delta < \varepsilon$. (See Fig. 6)

Therefore, (for $\delta < \varepsilon/4$) $\exists$ an edge $e \in G, e = y_i y_{i+1}$. From this we get the following upper bound for $d_G(x, y)$:

$$d_G(x, y) \leq \sum_0^n |y_i y_{i+1}| \leq \sum_0^n |x_i x_{i+1}| + 2\delta n$$

But $n < 2\text{length}(\gamma)/\varepsilon \leq 2\text{diam}(X)/\varepsilon$; therefore:

$$d_G(x, y) \leq |xy| + \delta \frac{4\text{diam}(X)}{\varepsilon} < |xy| + \varepsilon.$$

(because $\delta < \varepsilon^2/4\text{diam}(X)$).




FIGURE 6.

So, for any $\varepsilon > 0$, $\exists G = G_\varepsilon$ an ε -approximation of X . Then, $G_n \xrightarrow{\varepsilon} X$.

□

Corollary 2.52. *Let X be a compact length space. Then X is the Gromov-Hausdorff limit of a sequence $\{G_n\}_{n \geq 1}$ of finite graphs, isometrically embedded in X .*

Remark 2.53. (1) If $G_n \xrightarrow{\varepsilon} X$, $G_n = (V_n, E_n)$. If $\exists N_0 \in \mathbb{N}$ s.t.

$$(\star) \quad |E_n| \leq N_0, \forall n \in \mathbb{N},$$

then X is a finite graph.

(2) If condition $(\star)$ is replaced by:

$$(\star\star) \quad |V_n| \leq N_0, \forall n \in \mathbb{N},$$

then X will still be always a graph, but not necessarily finite(!)

3. THE EMBEDDING CURVATURE

3.1. Theoretical Setting. This is basically a *comparison-curvature* (as is the more "modern" CAT^{10} approach). This is done with quadruples instead of triangles (like in the Alexandrov-Topogonov method). It is in a sense a more natural idea, since quadruples are classically¹¹ the "minimal" geometric figures that allow the differentiation between metric spaces. This allows for a much more easier and rapid development of the theory than the triangle-based comparison. Moreover we shall show that the two Theories coincide on those metric space on which both can be applied, i.e. metric spaces that are (a) "planar" and (b) "rich enough" i.e. contain quadrangles, s.a. classical (PL-smooth) surfaces in $\mathbb{R}^3$.¹²

Definition 3.1. Let (M, d) be a metric space, and let $Q = \{p_1, \dots, p_4\} \subset M$, together with the mutual distances: $d_{ij} = d_{ji} = d(p_i, p_j)$; $1 \leq i, j \leq 4$. The set Q together with the set of distances $\{d_{ij}\}_{1 \leq i, j \leq 4}$ is called a *metric quadruple*.

Remark 3.2. One can define metric quadruples in slightly more abstract manner, without the aid of the ambient space: a metric quadruple being a 4 point metric space; i.e. $Q = (\{p_1, \dots, p_4\}, \{d_{ij}\})$, where the distances d_{ij} verify the axioms for a metric.

Before we proceed to the next definition, let us introduce the following

Notation S_κ denotes the complete, simply connected surface of constant curvature κ , i.e. $S_\kappa \equiv \mathbb{R}^2$, if $\kappa = 0$; $S_\kappa \equiv \mathbb{S}_{\sqrt{\kappa}}^2$, if $\kappa > 0$; and $S_\kappa \equiv \mathbb{H}_{\sqrt{-\kappa}}^2$, if $\kappa < 0$. Here $S_\kappa \equiv \mathbb{S}_{\sqrt{\kappa}}^2$ denotes the Sphere of radius $R = 1/\sqrt{\kappa}$, and $S_\kappa \equiv \mathbb{H}_{\sqrt{-\kappa}}^2$ stands for the Hyperbolic Plane of curvature $\sqrt{-\kappa}$, as represented by the Poincare Model of the plane disk of radius $R = 1/\sqrt{-\kappa}$

Definition 3.3. The *embedding curvature* $\kappa(Q)$ of the metric quadruple Q is defined be the curvature κ of S_κ into which Q can be isometrically embedded. (See Figures 7 and 8 for embeddings of the metric quadruple in S_κ and H_κ , respectively.)

We can now define the embedding curvature at a point in a natural way by passing to the limit (but without neglecting the existence conditions), more precisely:

Definition 3.4. Let (M, d) be a metric space, and let $p \in M$ be an accumulation point. Then p is said to have *Wald curvature* $\kappa_W(p)$ iff

- (i) $\nexists N \in \mathcal{N}(p)$, N linear¹³;
- (ii) $\forall \varepsilon > 0$, $\exists \delta > 0$ s.t. $Q = \{p_1, \dots, p_4\} \subset M$, and s.t. $d(p, p_i) < \delta$ ($i = 1, \dots, 4$)
 $\implies |\kappa(Q) - \kappa_W(p)| < \varepsilon$.

Remark 3.5. (1) If one uses the second (abstract) definition of the metric curvature of quadruples, then the very existence of $\kappa(Q)$ is not assured, as it is shown by the following

Counterexample 3.6. The metric quadruple of lengths

$$d_{12} = d_{13} = d_{14} = 1; d_{23} = d_{24} = d_{34} = 2$$

¹⁰ i.e. Cartan-Alexandrov-Topogonov

¹¹ as illustrated by the time-honored principles of Projective Geometry...

¹² In this sense CAT spaces are more "potent": they can be employed in studying mathematical objects that not (necessarilly) contain quadrangles, e.g. trees, Cayley graphs, etc..

¹³ The neighborhood N of p is called linear iff N is contained in a geodesic.




FIGURE 7.

admits no embedding curvature.

- (2) Even if a quadruple has an embedding curvature, it still may be not unique (even if Q is not linear), indeed, one can study the following examples:

Example 3.7. (a) The quadruple Q of distances $d_{ij} = \pi/2, 1 \leq i < j \leq 4$ is isometrically embeddable both in $S_0 = \mathbb{R}^2$ and in $S_1 = \mathbb{S}^2$.




FIGURE 8.

- (b) The quadruple Q of distances $d_{13} = d_{14} = d_{23} = d_{24} = \pi$, $d_{12} = d_{34} = 3\pi/2$ admits exactly two embedding curvatures: $\kappa_1 \in (1.5, 2)$ and $\kappa_2 = 3$. (See [BM].)

However, for "good" metric spaces¹⁴ the embedding curvature exists and it is unique. And, what is even more relevant for us, this embedding curvature coincides with the classical Gaussian curvature. The proof of this result is rather long and tedious, therefore we shall present here only a brief sketch of it. (This will prove to be somewhat redundant anyhow, in view of the more general results presented in the previous section, a fact but we shall emphasize later in our presentation).)

The Main ingredient for this proof, and for the analysis of yet another approach to curvature (the CAT one) is provided by the following string of propositions (which are just generalizations of the well known high-school triangle inequalities):

Proposition 3.8. *Let $\triangle(p_1, q_1, r_1) \subset \mathcal{S}_{\kappa_1}$ and $\triangle(p_2, q_2, r_2) \subset \mathcal{S}_{\kappa_2}$, s.t.*

$p_1q_1 = p_2q_2$, $p_1r_1 = p_2r_2$ and $\angle(q_1, p_1, r_1) = \angle(q_2, p_2, r_2)$.

Then: $\kappa_1 < \kappa_2 \implies q_1r_1 > q_2r_2$.

Proposition 3.9. *Let $p_1, q_1, r_1 \in \mathcal{S}_{\kappa_1}$, $p_2, q_2, r_2 \in \mathcal{S}_{\kappa_2}$ two isometric triples of points, s.t. the triple p_1, q_1, r_1 is not linear. Then:*

$\angle(q_1, p_1, r_1) < \angle(q_2, p_2, r_2)$, $\angle(p_1, q_1, r_1) < \angle(p_2, q_2, r_2)$, $\angle(q_1, r_1, p_1) < \angle(q_2, r_2, p_2)$.

Proposition 3.10. *Let $Q_1 = \{p_1, q_1, r_1, s_1\}$, $Q_2 = \{p_2, q_2, r_2, s_2\}$ be non-linear and non-degenerate quadruples in $\mathcal{S}_{\kappa_1}$, $\mathcal{S}_{\kappa_2}$, respectively. If $\triangle(p_1, q_1, r_1) \cong \triangle(p_2, q_2, r_2)$ and $\kappa_1 < \kappa_2$, then:*

- (1) $p_1s_1 = p_2s_2$, $q_1s_1 = q_2s_2 \implies r_1s_1 > r_2s_2$;
- (2) $r_1s_1 = r_2s_2$, $q_1s_1 = q_2s_2 \implies p_1s_1 > p_2s_2$;
- (3) $p_1s_1 = p_2s_2$, $r_1s_1 = r_2s_2 \implies q_1s_1 < q_2s_2$.

In order that we fully exploit the results above we need the following definition:

Definition 3.11. A metric quadruple $Q = Q(p_1, p_2, p_3, p_4)$, of distances $d_{ij} = \text{dist}(p_i, p_j)$, $i = 1, \dots, 4$, is called *semi-dependent* (a *sd-quad*, for brevity), iff 3 of its points are on a common geodesic, i.e. there exist 3 indices – e.g. 1,2,3 – s.t.: $d_{12} + d_{23} = d_{13}$.

Now we can easily formulate the following immediate consequence of Prop. 3.10.:

Corollary 3.12. *A sd-quad admits at most one embedding curvature.*

Unfortunately – as we have already noticed – in the general case the uniqueness of the embedding curvature is not guaranteed. However we can be a bit more explicit using the following definition:

Definition 3.13. Let $Q = \{p, q, r, s\}$ be a non-linear and non-degenerate quadruple. Q is called *planar* iff $\angle(q, p, r) + \angle(q, p, s) + \angle(s, p, r) = 2\pi$.

Then we have

Proposition 3.14. *Let $Q = \{p, q, r, s\}$ be a non-linear and non-degenerate quadruple in $\mathcal{S}_{\kappa}$. Then*

- (1) *If Q is planar, then it admits no isometric embedding in $\mathcal{S}_{\kappa_1}$, $\kappa_1 > \kappa$.*

¹⁴i.e. spaces that are locally "plane like"

(2) If Q is not planar, then it admits no isometric embedding in $\mathcal{S}_{\kappa_2}$, $\kappa_2 < \kappa$.

Corollary 3.15. *Let $Q = \{p, q, r, s\}$ be a non-linear and non-degenerate quadruple. Then Q has at most two different embedding curvatures.*

In fact we can state a much stronger assertion, of which Example 3.7.(a) is just a very particular case:

Proposition 3.16. $\forall p \in \mathcal{S}_\kappa$, and $\forall \kappa > 0$, $\exists U \in \mathcal{N}(p)$ s.t. $\exists$ a nonlinear, non-degenerate quadruple $Q \subset U$ of embedding curvature 0.

Proof. Let $\gamma_1, \gamma_2 \in U$, two great-circle arcs s.t. $\gamma_1 \cap \gamma_2 = p$.

Let $q_1, q_2 \in \gamma_1$ s.t. $pq_1 = pq_2 \neq 0$ and let $q \in \gamma_2$ s.t. $pq < \pi/2\sqrt{\kappa}$.¹⁵

Consider $\triangle(q'_1 q'_2, q') \subset \mathbb{R}^2$, $\triangle(q'_1 q'_2, q') \cong \triangle(q_1 q_2, q)$, let $p' = \frac{1}{2}q'_1 q'_2$, and let $h = q'p'$.




FIGURE 9.

Then since $0 < \kappa$, Proposition 3.10.(3) applied to the quadruples $\{q, q_1, q_2, p\}$ and $\{q', q'_1, q'_2, p'\}$ implies that $h < pq$.

Now let $x \in \gamma_2$, x between p and q , and let $x' \in \mathbb{R}^2$ s.t. $\triangle(q'_1 q'_2, x') \cong \triangle(q_1 q_2, x)$ s.t. x and q' are on different sides of the line $\overleftrightarrow{q_1 q_2}$. Then, $x = p \Rightarrow xq > x'q'$, and $x = q \Rightarrow xq = 0 < x'q' = 2h$, where, in this case $x' = q''$. (See Figure 9.)

¹⁵i.e a quarter of the length of a great circle in $\mathcal{S}_\kappa$

Then it follows from a continuity argument that $\exists x_0 \in \gamma_2$, x_0 between p and q , s.t. $x_0q = x'_0q'$, thus implying that $\{q_1, q_2, q, x\} \cong \{q'_1, q'_2, q', x'\}$. $\square$

Remark 3.17. $\{q_1, q_2, q, x\}$ is planar, while $\{q'_1, q'_2, q', x'\}$ is not planar.

3.2. The Wald Curvature vs. Gauss Curvature. The discussion above would be nothing more than a nice intellectual exercise were it not for the fact that the metric (Wald) and the classical (Gauss) curvatures coincide whenever the second notion makes sense, that is for smooth (i.e. of class $\geq C^2$) surfaces in $\mathbb{R}^3$. More precisely the following theorem holds:

Theorem 3.18. (Wald) *Let $S \subset \mathbb{R}^3$, $S \in C^m$, $m \geq 2$ be a smooth surface. Then $\kappa_W(p)$ exists, for all $p \in S$, and $\kappa_W(p) = \kappa_G(p)$, $\forall p \in M$.*

Moreover, Wald also proved that a partial reciprocal theorem holds, more precisely he proved the following:

Theorem 3.19. *Let M be a compact and convex metric space. If $\kappa_W(p)$ exists, for all $p \in M$, then M is a smooth surface and $\kappa_W(p) = \kappa_G(p)$, $\forall p \in M$.*

Remark 3.20. If one tries to restrict oneself, in the building of Definition 3.4. only to sd-quads, then Theorem 3.19. holds only if the following presumption is added:

Condition 3.21. *M is locally homeomorphic to $\mathbb{R}^2$.*

However the proof of this facts is involved and, as such, beyond the scope of this presentation. Therefore we shall restrict ourselves to a succinct description of the principal steps towards the proofs.

The basic idea is to show that if a metric M space admits a Wald curvature at any point, then M is locally homeomorphic to $\mathbb{R}^2$, thus any metric proprieties of $\mathbb{R}^2$ can be translated to M , (in particular the first fundamental form).

The first of these partial results is:

Theorem 3.22. *Let M be a convex metric space. Then M admits at most one Wald curvature $\kappa_W(p)$, $\forall p \in M$.*

Proof By Corollary 3.12. it suffices to prove that any disk neighborhood $B(p; \rho) \in \mathcal{N}(p)$ contains a non degenerate sd-quad.

Without loss of generality one can assume that $B(p; \rho)$ contains three points p_1, p_2, p_3 s.t. $d(p, p_i) < \rho/2$, $i = 1, 2, 3$.¹⁶ Then, by the convexity of M it follows that $\exists q \in M$ s.t. $p \neq p_2, p_3$ and $p_2q + p_3q = p_2p_3$. But $p_2p_3 \leq pp_2 + pp_3 < \rho \implies (pq < \rho/2) \vee (pp_2 < \rho/2)$. In the first inequality holds, then $pq \leq pp_2 + p_2q < \rho$, i.e. $q \in B(p; \rho)$; and if the second one holds, then $pd \leq pp_3 + p_3q < \rho$, i.e. $q \in B(p; \rho)$. But $p \neq q$, therefore p, p_2, p_3, q are not linear. $\square$

Our next step will be to analyze those neighborhoods that display "a normal behavior", both metrically and curvature-wise: that is precisely those disk neighborhoods in which the Wald curvature is defined and ranges over a small, bounded set of values prescribed by the very radius of the disk:

Definition 3.23. A disk neighborhood $B(p; \rho)$, $\rho > 0$ is called *regular* iff $\forall$ non-degenerate quadruple $Q \subset B(p; \rho)$, $\kappa_W(Q)$ exists and $|\kappa_W(Q)| < \pi^2/16\rho^2$.

¹⁶See [B]

Remark 3.24. If $\kappa_W(p)$ exists, then for any sufficiently small ρ , $B(p; \rho)$ will be regular.

It turns out that regular neighborhoods, in *compact, convex* spaces have the following "nice" (i.e. Real Plane like) proprieties:

Proposition 3.25. *Let M be a compact, convex metric space and let $B(p; \rho) \subset M$ be a regular neighborhood. Then if a non-degenerate quadruple $Q \subset B(p; \rho)$ contains two linear triples of points, then Q is linear.*

Proposition 3.26. *Let M be a compact, convex metric space. Then:
 $\forall p \in M$ and $\forall B(p; \rho)$ regular, $\exists q, r \in B(p; \rho)$ s.t. p, q, r are not linear.*

Proposition 3.27. *Any regular neighborhood $B(p; \rho)$ of a compact, convex metric space is strictly convex, i.e. $q, r \in B(p; \rho) \implies \text{int}(qr) \subset B(p; \rho)$.*

While the proof of this last Proposition is lengthy, that of the following important Corollary is not:

Corollary 3.28. *Let $B(p; \rho)$ be a regular neighborhood. Then, $\forall q, r \in B(p; \rho)$, $\exists! qr$ and $\text{int}(qr) \subset B(p; \rho)$.*

Proof. By the convexity of $B(p; \rho)$ it follows the existence of at least one geodesic qr , $\forall q, r \in B(p; \rho)$. If $s \in \text{int}(qr)$, then by the proposition above we have that $s \in B(p; \rho)$. It follows that $B(p; \rho)$ contains all the geodesics with end points q, r . Hence, by Proposition 3.25., the geodesic segment qr is unique. $\square$

We can now begin to prove that a compact, convex metric space locally mimics $\mathbb{R}^2$. We start by showing that the *sinus* function is defined on M , thus allowing for angle measure (hence for the definition of Polar Coordinates on regular neighbourhoods¹⁷).

First, let M be as before, and let $p \in M$ s.t. $\kappa_W p$ exists. Let $q, r \in B(p; \rho)$, $q \neq p \neq r$, where $B(p; \rho)$ is a regular neighborhood of p . Then, $\forall x \in [0, \min\{pq, pr\})$, define $q(x) \in pq$, $r(x) \in pr$ by: $d(p, q(x)) = x = d(p, r(x))$, and let $d(x) = d(q(x), r(x))$ (see Figure 10 below).

Proposition 3.29. *The following limit exists:*

$$\lim_{x \rightarrow 0} \frac{d(x)}{x}.$$

We omit the proof since it is rather involved (but canonical for any axiomatic approach to Euclidian Geometry – see, for instance, [B], [RR].)

Now we can define the measure of angles at p :

Definition 3.30. The measure of the angle $\angle(q, p, r)$ is given by:

$$m(\angle(q, p, r)) \triangleq 2\arcsin\left(\frac{1}{2} \lim_{x \rightarrow 0} \frac{d(x)}{x}\right).$$

Remark 3.31. The Definition above enables us to define Polar Coordinates on regular neighborhoods¹⁸ in the following manner:

Let $p_1, p_2 \in B(p; \rho)$ s.t. p, p_1, p_2 are not collinear. (Such points exist by Proposition 3.26.). To every point $q \in B(p; \rho)$ we associate the following pair of real

¹⁷ In the same way geodesic polar coordinates are used on classical surfaces.

¹⁸ ... and once Coordinates (be they Polar or Cartesian) are introduced, the (local) homomorphism with $\mathbb{R}^2$ is immediate.




FIGURE 10.

numbers (defining the Polar Coordinates of q relative to the frame determined by p, p_1, p_2): $(r(q), \theta(q))$, where

$$r(q) \triangleq d(p, q)$$

and

$$\theta(q) \triangleq \begin{cases} m(\angle(q, p, p_1)) & \text{if } |m(\angle(p_2, p, p_1)) - m(\angle(q, p, p_1))| = m(\angle(q, p, p_1)) ; \\ 2\pi - m(\angle(q, p, p_1)) & \text{if } |m(\angle(p_2, p, p_1)) - m(\angle(q, p, p_1))| \neq m(\angle(q, p, p_1)) . \end{cases}$$

We can now safely state the foretold homomorphism result:

Proposition 3.32. *Any convex, compact metric space is locally homeomorphic to the real plane.*

4. COMPUTING EMBEDDING CURVATURE

In this section we develop formulas for the computation of Embedding Curvature of Quadruples. First we follow the classical approach of Wald-Blumenthal that employs the so-called *Cayley-Menger determinants* (see below). Unfortunately, the formulas obtained, albeit precise are transcendental. Therefore we present, in the next subsection, the approximate formulas developed by C.V. Robinson.

4.1. Embedding Curvature – The Determinant Approach. Given a general metric quadruple $Q = Q(p_1, p_2, p_3, p_4)$, of distances $d_{ij} = \text{dist}(p_i, p_j)$, $i = 1, \dots, 4$, we denote by $D(Q) = D(p_1, p_2, p_3, p_4)$ the following determinant:

$$(4.1) \quad D(p_1, p_2, p_3, p_4) = \begin{vmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & d_{12}^2 & d_{13}^2 & d_{14}^2 \\ 1 & d_{12}^2 & 0 & d_{23}^2 & d_{24}^2 \\ 1 & d_{13}^2 & d_{23}^2 & 0 & d_{34}^2 \\ 1 & d_{14}^2 & d_{24}^2 & d_{34}^2 & 0 \end{vmatrix}$$

Then the *embedding curvature* $\kappa(Q)$ of Q is given – depending upon the embedding space (i.e. upon the sign of the curvature) – by the following formulae:

$$(4.2) \quad \kappa(Q) = \begin{cases} 0 & \text{if } D(Q) = 0; \\ \kappa, \kappa < 0 & \text{if } \det(\cosh \sqrt{-\kappa} \cdot d_{ij}) = 0; \\ \kappa, \kappa > 0 & \text{if } \det(\cos \sqrt{\kappa} \cdot d_{ij}) \text{ and } \sqrt{\kappa} \cdot d_{ij} \leq \pi \\ & \text{and all the principal minors of order 3 are } \geq 0. \end{cases}$$

The determinant $D(Q) = D(p_1, p_2, p_3, p_4)$ is called the *Cayley-Menger determinant* (of the points $p_1, \dots, p_4$)¹⁹ and, in order to prove (4.2) we need first to investigate some of its properties.

We start with the following

Lemma 4.1. *Let: $p_1, \dots, p_4$ be points in $\mathbb{R}^3$. Then:*

$$(4.3) \quad \text{Gram}(\overrightarrow{p_1 p_2}, \overrightarrow{p_1 p_3}, \overrightarrow{p_1 p_4}) = \frac{1}{8} D(p_1, p_2, p_3, p_4);$$

where

$$\text{Gram}(\overrightarrow{p_1 p_2}, \overrightarrow{p_1 p_3}, \overrightarrow{p_1 p_4}) \stackrel{\text{def}}{=} \det(\overrightarrow{p_1 p_i} \cdot \overrightarrow{p_1 p_j})_{i,j=2,3,4}$$

(Here “.” denotes the standard scalar (dot) product in $\mathbb{R}^3$.)

Proof. Use expansion and manipulation of determinants. □

Since is a known fact that:

$$\text{Gram}(\overrightarrow{p_1 p_2}, \overrightarrow{p_1 p_3}, \overrightarrow{p_1 p_4}) = (\text{Vol}(p_1, p_2, p_3, p_4))^2;$$

where $\text{Vol}(p_1, p_2, p_3, p_4)$ denotes the (un-oriented) volume of the parallelepiped determined by the vertices $p_1, \dots, p_4$ (and with edges $\overrightarrow{p_1 p_2}, \overrightarrow{p_1 p_3}, \overrightarrow{p_1 p_4}$); formula (2.3) shows that:

$$(4.4) \quad D(p_1, p_2, p_3, p_4) = 8(\text{Vol}(p_1, p_2, p_3, p_4))^2.$$

Therefore the following assertion is immediate:

Proposition 4.2. *The points $p_1, \dots, p_4$ are the vertices of a simplex in $\mathbb{R}^3$ iff $D(p_1, p_2, p_3, p_4) \neq 0$:*

However, we can prove the much strong result bellow:

Theorem 4.3. *Let $d_{ij} > 0, 1 \leq 4, i \neq j$.*

Then there exists a simplex $T = T(p_1, \dots, p_4) \subseteq \mathbb{R}^3$ s.t. $\text{dist}(x_i, x_j) = d_{ij}, i \neq j$; iff

$D(p_i, p_j) < 0, (\forall) \{i, j\} \subset \{1, \dots, 4\}$ and

$D(p_i, p_j, p_k) > 0, (\forall) \{i, j, k\} \subset \{1, \dots, 4\}$;

where, for instance,

$$D(p_1, p_2) = \begin{vmatrix} 0 & 1 & 1 \\ 1 & 0 & d_{12}^2 \\ 1 & d_{12}^2 & 0 \end{vmatrix}$$

and

$$D(p_1, p_2, p_3) = \begin{vmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & d_{12}^2 & d_{13}^2 \\ 1 & d_{12}^2 & 0 & d_{23}^2 \\ 1 & d_{13}^2 & d_{23}^2 & 0 \end{vmatrix};$$

¹⁹This definition readily generalizes to any dimension, as do the results bellow.

etc...

In fact, the necessary and sufficient condition above can be relaxed, indeed one can also show that the following holds²⁰:

Proposition 4.4.²¹ *Let $d_{ij} > 0, 1 \leq i, j \leq 4, i \neq j$.*

Then there exists a simplex $T = T(p_1, \dots, p_4) \subseteq \mathbb{R}^3$ s.t. $\text{dist}(x_i, x_j) = d_{ij}, i \neq j$; iff $D(p_1, p_2, p_3, p_4) \neq 0$ and $\text{sign } D(p_1, p_2, p_3, p_4) = +1$.

Proof.(Sketch) Sufficient to show (by using standard operations on determinants) that:

$$D(p_1, p_2, p_3)D(p_1, \dots, \hat{p}_i, \dots, p_4) = M_{i4}^2 + D(p_1, \dots, \hat{p}_i, \dots, p_4)D(p_1, p_2, p_3, p_4);$$

where M_{i4} is the cofactor (in D) of d_{i4}^2 , and were we used the notation:

$$\{p_1, \dots, \hat{p}_i, \dots, p_4\} = \{p_1, p_2, p_3, p_4\} \setminus \{p_i\}.$$

□

Proving the formula for the spherical and hyperbolic cases would prove to be too technical for this limited exposition; suffice to say that they essentially reproduce the proof given in the Euclidian case, and tacking into account the fact that performing computations in the spherical (resp. hyperbolic) metric one has to replace the distances d_{ij} by $\cos d_{ij}$ (resp. $\cosh d_{ij}$)²².

4.2. Embedding Curvature – Approximate Formulas. The formulas we just developed in are not only transcendental, but also the computed curvature may fail to be unique (see preceding section). However, uniqueness is guaranteed for sd-quads. Moreover, the relatively simple geometric setting of sd-quads allows for the development of simple (i.e. rational) formulas for the approximation of the Embedding Curvature.

Proposition 4.5. *Given the metric quadruple $Q = Q(p_1, p_2, p_3, p_4)$, of distances $d_{ij} = \text{dist}(p_i, p_j), i = 1, \dots, 4$, the embedding curvature $\kappa(Q)$ is well approximated by:*

$$(4.5) \quad K(Q) = \frac{6(\cos \angle_0 2 + \cos \angle_0 2')}{d_{24}(d_{12} \sin^2(\angle_0 2) + d_{23} \sin^2(\angle_0 2'))}$$

where: $\angle_0 2 = \angle(p_1 p_2 p_4), \angle_0 2' = \angle(p_3 p_2 p_4)$ represent the angles of the Euclidian triangles of sides d_{12}, d_{14}, d_{24} and d_{23}, d_{24}, d_{34} , respectively.

The error R can be estimated by using the following inequality:

$$(4.6) \quad |R| = |R(Q)| = |\kappa(Q) - K(Q)| < 4\kappa^2(Q) \text{diam}^2(Q) / \lambda(Q)$$

where we put: $\lambda(Q) = d_{24}(d_{12} \sin \angle_0 2 + d_{23} \sin \angle_0 2') / S^2$, and where $S = \text{Max}\{p, p'\}; 2p = d_{12} + d_{14} + d_{24}, 2p' = d_{32} + d_{34} + d_{24}$.

Proof The basic idea of the proof is to recreate, in a general metric setting, the Gauss Map – in this case one measures the curvature by the amount of "bending" one has to apply to a general planar quadruple so that it may be "straightened" (i.e. isometrically embedded as a *sd-quad*) in some S_κ .

Consider two plane²³ triangles $\triangle p_1 p_2 p_4$ and $\triangle p_2 p_3 p_4$, and denote by $\triangle p_{1,k} p_{2,k} p_{4,k}$

²⁰ For the direct proof of Theorem 4.3., see [Be] or, alternatively [B]

²¹ We formulate this result – for convenience and practicality – for the case $n = 3$, only. However it is readily generalized to any dimension.

²² See [B] for the full details

²³ i.e. embedded in $R^2 \equiv S_0$

and $\triangle p_{2,k}p_{3,k}p_{4,k}$ their respective isometric embeddings into $\mathcal{S}_k$. Then $p_{i,k}p_{j,k}$ will denote the geodesic (of $\mathcal{S}_k$) through $p_{i,k}$ and $p_{j,k}$. Also, let $\angle_k 2$ and $\angle_k 2'$ denote, respectively, the following angles of $\triangle p_{1,k}p_{2,k}p_{4,k}$ and $\triangle p_{2,k}p_{3,k}p_{4,k}$: $\angle_k 2 = \angle p_{1,k}p_{2,k}p_{4,k}$ and $\angle_k 2' = \angle p_{2,k}p_{3,k}p_{4,k}$. (See Fig. 11)




FIGURE 11.

But $\angle_k 2$ and $\angle_k 2'$ are strictly increasing as functions of k . Therefore the equation

$$(4.7) \quad \angle_k 2 + \angle_k 2' = \pi$$

has at most one solution k^* , i.e. k^* represents the unique value for which the points p_1, p_2, p_4 are on a geodesic in $\mathcal{S}_k$ (for instance on p_1p_4).

But that means that k^* is precisely the Embedding Curvature, i.e. $k^* = \kappa(Q)$, where $Q = Q(p_1, p_2, p_3, p_4)$.

Equation (4.7) is equivalent to

$$\cos^2 \frac{\angle_{k^*} 2}{2} + \cos^2 \frac{\angle_{k^*} 2'}{2} = 1$$

The basic idea being the comparison between metric triangles with equal sides, embedded in $\mathcal{S}_0$ and $\mathcal{S}_k$, respectively, it is natural to consider instead of the previous equation, the following:

$$(4.8) \quad \theta(k, 2) \cdot \cos^2 \frac{\angle_0 2}{2} + \theta(k, 2') \cdot \cos^2 \frac{\angle_0 2'}{2} = 1$$

where we denote:

$$\theta(k, 2) := \frac{\cos^2 \frac{\angle_{k^*} 2}{2}}{\cos^2 \frac{\angle_0 2}{2}}; \quad \theta(k, 2') := \frac{\cos^2 \frac{\angle_{k^*} 2'}{2}}{\cos^2 \frac{\angle_0 2'}{2}}.$$

Since we want to approximate $\kappa(Q)$ by $K(Q)$ we shall resort – naturally – to expansion into MacLaurin series. We are able to do this because of the existence of the following classical formulas:

$$\cos^2 \frac{\angle_k 2}{2} = \frac{\sin(p\sqrt{k}) \cdot \sin(d\sqrt{k})}{\sin(d_{12}\sqrt{k}) \cdot \sin(d_{24}\sqrt{k})}; \quad k > 0;$$

$$\cos^2 \frac{\angle_k 2}{2} = \frac{\sinh(p\sqrt{k}) \cdot \sinh(d\sqrt{k})}{\sinh(d_{12}\sqrt{k}) \cdot \sinh(d_{24}\sqrt{k})}; \quad k < 0;$$

and, of course

$$\cos^2 \frac{\angle_0 2}{2} = \frac{pd}{d_{12}d_{24}};$$

were: $d = p - d_{14} = (d_{12} + d_{24} - d_{14})/2$.²⁴

By using the development into series of $f_1(x) = \frac{\sin \sqrt{x}}{\sqrt{x}}$ and $f_2(x) = \frac{\sinh \sqrt{x}}{\sqrt{x}}$; one (easily) gets the desired expansion for $\theta(k, 2)$:

$$(4.9) \quad \theta(k, 2) = 1 + \frac{1}{6}kd_{12}d_{24}(\cos(\angle_0 2) - 1) + r;$$

where: $|r| < \frac{3}{8}k^2p^4$, for $|kp^2| < 1/16$. By applying (4.9.) to (4.8), we receive:

$$(4.10) \quad [1 + \frac{1}{6}k^*d_{12}d_{24}(\cos(\angle_0 2) - 1) + r] \cos^2 \frac{\angle_0 2}{2} + \\ [1 + \frac{1}{6}k^*d_{23}d_{24}(\cos(\angle_0 2') - 1) + r'] \cos^2 \frac{\angle_0 2'}{2} = 1;$$

for: $|r| + |r'| < \frac{3}{4}(k^*)^2(Max\{p, p'\})^4 = \frac{3}{4}(k^*)^2S^4$.

By solving linear equation (in variable k^*) (4.10) and using some elementary trigonometric transformation one has:

$$k^* = \frac{6(\cos \angle_0 2 + \cos \angle_0 2')}{d_{24}(d_{12} \sin^2(\angle_0 2) + d_{23} \sin^2(\angle_0 2'))} + R$$

where:

$$|R| < \frac{12(|r| + |r'|)}{d_{24}(d_{12} \sin^2(\angle_0 2) + d_{23} \sin^2(\angle_0 2'))} < \frac{9(k^*)^2 \max\{p, p'\}}{d_{24}(d_{12} \sin^2(\angle_0 2) + d_{23} \sin^2(\angle_0 2'))}$$

But $k^* \equiv \kappa(Q)$ so we get the desired formula (4.5).

To prove the correctness of the bound (4.6) one has only to observe that:

$$S = Max\{p, p'\} < 2diam(Q), \quad (diam(Q) = \max_{1 \leq i < j \leq 4} \{d_{ij}\}),$$

and perform the necessary arithmetic manipulations. □

Remark 4.6. (a) The function $\lambda = \lambda(Q)$ is continuous and 0-homogenous as a function of the d_{ij} -s. Moreover: $\lambda(Q) \geq 0$ and $\lambda(Q) = 0 \Leftrightarrow \sin \angle_0 2 = \sin \angle_0 2' = 0$, i.e. iff Q is linear. [Therefore for sd-quads $\lambda(Q) > 0$ and, moreover, $\lambda(Q) \rightarrow 0 \Rightarrow Q \rightarrow \text{linearity}$.]

(b) Since $\lambda(Q) \neq 0$ it follows that: $K(Q) \in \mathbb{R}$ for any quadrangle Q . In addition: $sign(k(Q)) = sign(K(Q))$.

(c) If Q is any sd-quad, then $\kappa^2(Q)diam^2(Q)/\lambda(Q) < \infty$. Moreover, if $\lambda(Q) \not\gg 0$,²⁵ then $\kappa^2(Q)diam^2(Q)/\lambda(Q) \not\gg 0$ i.e. $|R|$ is *small* if Q is not close to linearity.

In this case $|R(Q)| \sim diam^2(Q)$ (for any given Q).

²⁴and the analogous formulas for $\cos^2 \frac{\angle_{k'} 2}{2}$.

²⁵i.e. for not very small values of $\lambda(Q)$

Since the Gaussian curvature $k_G(p)$ at a point p is given by:

$$k_G(p) = \lim_{n \rightarrow 0} \kappa(Q_n);$$

where $Q_n \rightarrow Q = \square p_1 p p_3 p_4$; $\text{diam}(Q_n) \rightarrow 0$, from Remark 4.6.(c) we immediately infer that the following holds²⁶:

Theorem 4.7. *Let S be a differentiable surface. Then, for any point $p \in S$:*

$$k_G(p) = \lim_{n \rightarrow 0} K(Q_n);$$

for any sequence $\{Q_n\}$ of sd-quads that satisfy the following condition:

$$Q_n \rightarrow Q = \square p_1 p p_3 p_4; \text{diam}(Q_n) \rightarrow 0.$$

Remark 4.8. In the following special cases even "nicer" formulas are obtained:

(1) If $d_{12} = d_{32}$, then

$$(4.11) \quad K(Q) = \frac{12}{d_{13} \cdot d_{24}} \cdot \frac{\cos \angle_0 2 + \cos \angle_0 2'}{\sin^2 \angle_0 2 + \sin^2 \angle_0 2'};$$

(here we have of course: $d_{13} = 2d_{12} = 2d_{32}$); or, expressed as a function of distances alone:

$$(4.12) \quad K(Q) = 12 \frac{2d_{12}^2 + 2d_{24}^2 - d_{14}^2 - d_{13}^2}{8d_{12}^2 d_{24}^2 - (d_{12}^2 + d_{24}^2 - d_{14}^2)^2 - (d_{12}^2 + d_{24}^2 - d_{34}^2)^2}$$

(2) If $d_{12} = d_{32} = d_{24}$ and if the following condition also holds:

(3) $\angle_0 2' = \pi/2$; i.e. if $d_{34}^2 = d_{12}^2 + d_{24}^2$ or, considering (2), also: $d_{34}^2 = 2d_{12}^2$ then

$$(4.13) \quad K(Q) = \frac{6 \cos \angle_0 2}{d_{12}(1 + \sin^2 \angle_0 2)} = \frac{2d_{12}^2 - d_{14}^2}{4d_{12}^4 + 4d_{14}^2 d_{12}^2 - d_{14}^4}.$$

5. APPENDIX 1 – THE Menger AND HAANTJES CURVATURES

Better known than the Wald Curvature, the Menger Curvature is a metric definition of curvature of *curves*, as is the Haantjes Curvature. As such they can be employed as *sectional curvatures* to approximate curvature of triangulated surfaces. We begin by introducing the the Menger Curvature: this is a metric expression for the circum-radius of a triangle²⁷, based upon elementary high-school formulas:

Definition 5.1. Let (M, d) be a metric space, and let $p, q, r \in M$ be three distinct points. Then:

$$K_M(p, q, r) = \frac{\sqrt{(pq + qr + rp)(pq + qr - rp)(pq - qr + rp)(-pq + qr + rp)}}{pq \cdot qr \cdot rp};$$

is called the *Menger Curvature* of the points p, q, r .

We can now define the Menger Curvature at a given point by passing to the limit:

Definition 5.2. Let (M, d) be a metric space and let $p \in M$ be an accumulation point. Then M has at p *Menger Curvature* $\kappa_M(p)$ iff
 $\forall \varepsilon > 0, \exists \delta > 0$ s.t. $d(p, p_i) < \delta$; $i = 1, 2, 3 \implies |K(Q) - \kappa_M(p)| < \varepsilon$.

²⁶and gives theoretical justification to the algorithm

²⁷thus giving in the limit a metric definition of the Osculatory Circle

Remark 5.3. The apparent equivalent notion of *Alt Curvature*, in which one uses only two points converging to the third, is in fact more general, where we define the *Arp curvature* by:

Definition 5.4. Let (M, d) be a metric space and let $P \in M$ be an accumulation point. Then M has at p *Alt Curvature* $\kappa_A(p)$ iff the following limit exists

$$\kappa_A(p) \triangleq \lim_{q, r \rightarrow p} K(p, q, r).$$

However, both $\kappa_M(p)$ and $\kappa_A(p)$ suffer from the same imperfection: since they are both modelled closely after the Euclidian Plane, they convey this Euclidian type of curvature upon the space they are defined on. However, the next definition doesn't mimic closely $\mathbb{R}^2$ so it better fitted for generalizations:

Definition 5.5. Let (M, d) be a metric space and let $c : I = [0, 1] \xrightarrow{\sim} M$ be a homeomorphism, and let $p, q, r \in c(I)$, $q, r \neq p$. Denote by $\widehat{qr}$ the arc of $c(I)$ between q and r , and by qr segment from q to r . (See Figure 12 below.)




FIGURE 12.

Then c has *Haantjes Curvature* $\kappa_H(p)$ at the point p iff:

$$\kappa_H^2(p) = 24 \lim_{q, r \rightarrow p} \frac{l(\widehat{qr}) - d(q, r)}{(l(\widehat{qr}))^3};$$

where " $l(\widehat{qr})$ " denotes the length²⁸ of $\widehat{qr}$.

Remark 5.6. κ_H exists only for rectifiable curves, but if κ_M exists at any point p of c , then c is rectifiable.

Remark 5.7. Evidently we have the following relationship between curvatures:

$$\exists \kappa_M \implies \exists \kappa_A.$$

while

$$\exists \kappa_A \not\Rightarrow \exists \kappa_M.$$

However, we can prove the following theorem:

²⁸ given by the intrinsic metric induced by d

Theorem 5.8. *Let $c : I \rightarrow M$ be a rectifiable curve, and let $p \in M$. If κ_A (or κ_M) exists, then $\kappa_H(p)$ exists and*

$$\kappa_A = \kappa_H(p).$$

Remark 5.9. This last result and the Remark preceding it allow us to employ any of the curvatures above in estimating curvatures of smooth curves on triangulated surfaces.

6. APPENDIX 2 – THE RINOW CURVATURE

The curvatures introduced before may seem a bit archaic in comparison to the more fashionable approach of *comparison triangles*, with their ar reaching applications. We present here one of these comparison criteria and show its equivalence with the Wald curvature. We start with the following definition:

Definition 6.1. Let (M, d) be a metric space, together with the intrinsic metric induced by d . Let $R = \text{int}(R) \subseteq M$ be a region of M . We say that R is a *region of curvature $\leq \kappa$* ($\kappa \in \mathbb{R}$) iff

- (1) $\forall p, q \in R, \exists$ a geodesic segment $pq \subset R$;
- (2) $\forall T(p, q, r) \subset R$ is isometrically embeddable in $\mathcal{S}_\kappa$;
- (3) If $T(p, q, r) \subset R$ and $x \in pq, y \in pr$, and if the points $p_\kappa, q_\kappa, r_\kappa, x_\kappa, y_\kappa \in \mathcal{S}_\kappa$ satisfy the following conditions:
 - (a) $T(p, q, r) \cong T(p_\kappa, q_\kappa, r_\kappa)$;
 - (b) $T(p, q, x) \cong T(p_\kappa, q_\kappa, x_\kappa)$;
 - (c) $T(p, r, y) \cong T(p_\kappa, r_\kappa, y_\kappa)$;
then $xy \leq x_\kappa y_\kappa$.

By replacing the condition: " $xy \leq x_\kappa y_\kappa$ " with: " $xy \geq x_\kappa y_\kappa$ ", we obtain the definition of a *region of curvature $\geq \kappa$* . (See Fig. 13.)




FIGURE 13.

We now pass to the localization of the Definition above:

Definition 6.2. Let (M, d) be a metric space, together with the intrinsic metric induced by d , and let $p \in M$ be an accumulation point. Then M has at p *Rinow Curvature* $\kappa_R(p)$ iff

- (i) $\nexists N \in \mathcal{N}(p)$, N linear;
- (ii) $\forall \varepsilon > 0$, $\exists \delta.0$, s.t. $B(p; \delta)$ is (a) a region of Rinow curvature $\leq \kappa_R(p) + \varepsilon$ and (b) a region of Rinow curvature $\geq \kappa_R(p) - \varepsilon$.

While its greater generality endows the Rinow curvature with more flexibility in applications and makes it easier in generalization, it is even more difficult to compute than Wald Curvature. However this quandary was has an almost ideal solution, due to Kirk (see [K]), solution which we briefly expose here:

Definition 6.3. Let M be a compact, convex metric space, and let $p \in M$. If $\kappa_W(p)$ exists, then $\kappa_R(p)$ exists, and $\kappa_R(p) = \kappa_W(p)$.

Unfortunately, since $\kappa_R(p)$ makes no presumption of dimensionality, the existence of $\kappa_R(p)$ does not imply the existence of $\kappa_W(p)$.

Counterexample 6.4. Let $M \equiv \mathbb{R}^3$. Then $\kappa_R(p) \equiv 0$ but $\kappa_W(p)$ does not exist at any point, since every neighborhood contains linear quadruples.

The solution (due to Kirk) of this problem is to consider the *Modified Wald curvature* κ_{WK} , defined as follows:

Definition 6.5. Let (M, d) be a metric space, together with the intrinsic metric induced by d , and let $p \in M$. Then M has at p *Modified Wald curvature Curvature* $\kappa_{WK}(p)$ iff

- (i) $\nexists N \in \mathcal{N}(p)$, N linear;
- (ii) $\forall \varepsilon > 0$, $\exists \delta.0$, s.t. if $Q \subset B(p; \delta)$ is a non-degenerate sd-quad, then $\kappa_W(Q)$ exists and $|\kappa_{WK}(p) - \kappa_W(Q)| < \varepsilon$.

Remark 6.6. $\exists \kappa_W(p) \implies \exists \kappa_{WK}(p)$ but $\exists \kappa_{KW}(p) \not\Rightarrow \kappa_W(p)$.

This modified curvature indeed represents the wished for solution, as proved by the following to Theorems:

Theorem 6.7. Let (M, d) be a metric space. Then:

$$\exists \kappa_R(p) \implies \exists \kappa_{WK}(p) \text{ and } \kappa_R(p) = \kappa_{WK}(p).$$

Theorem 6.8. Let (M, d) be a metric space together with the associated intrinsic metric, and let $p \in M$. Then, if

(i) $\kappa_{WK}(p)$;

and if

(ii) $\exists B(p; \rho) \in \mathcal{N}(p)$, s.t. $qr \subset B(p; \rho)$, $\forall q, r \in B(p; \rho)$;

then $\kappa_R(p)$ exists and $\kappa_R(p) = \kappa_{WK}(p)$.

7. APPENDIX 3 – THE RADIUS FORMULA

The Cayley-Menger determinant allows one to express not only the volume and area²⁹ of simplices in $\mathbb{R}^n$ but (as expected) it may be used to compute the radius

²⁹The 2-dimensional analogue of Formula (4.4) for the area of the triangle $T(p_1, p_2, p_3)$ being:

$$(\text{Area}(p_1, p_2, p_3))^2 = -D(p_1, p_2, p_3).$$

of the circumscribed sphere around an Euclidian simplex. To be more precise, we have the following result³⁰:

Theorem 7.1. (1) *The radius $R = R(p_1, p_2, p_3, p_4)$ of the sphere circumscribed around the tetrahedron $T(p_1, p_2, p_3, p_4) \in \mathbb{R}^3$ is given by:*

$$R^2 = -\frac{1}{2} \frac{\Delta(p_1, p_2, p_3, p_4)}{D(p_1, p_2, p_3, p_4)}$$

where:

$$\Delta(p_1, p_2, p_3, p_4) = \begin{vmatrix} 0 & d_{12}^2 & d_{13}^2 & d_{14}^2 \\ d_{12}^2 & 0 & d_{23}^2 & d_{24}^2 \\ d_{13}^2 & d_{23}^2 & 0 & d_{34}^2 \\ d_{14}^2 & d_{24}^2 & d_{34}^2 & 0 \end{vmatrix}$$

(2) *The points $p_1, p_2, p_3, p_4, p_5 \in \mathbb{R}^3$ are coplanar or co-spherical iff*

$$\Delta(p_1, p_2, p_3, p_4, p_5) = 0.$$

Proof

(1) If $p_0 \in \mathbb{R}^3$ is s.t. $d_{0i} = R$, $i = 1, \dots, 5$, then by direct computation we obtain:

$$\Gamma(p_0, \dots, p_5) = -2R^2\Gamma(p_1, \dots, p_5) - \Delta(p_1, p_2, p_3, p_4, p_5);$$

from which the desired formula follows immediately if we chose p_0 as the center of the sphere circumscribed around the points p_1, p_2, p_3, p_4, p_5 .

(2) ($\implies$) Let $\{x^1, x^2, x^3\}$ be any orthonormal coordinate frame for $\mathbb{R}^3$, and let p_i^j ; $i = 1, \dots, 5$; $j = 1, 2, 3$; represent the coordinates of the points p_1, p_2, p_3, p_4, p_5 relative to this coordinate system. Then p_1, p_2, p_3, p_4, p_5 belong to the same sphere or plane iff $\exists (a, b, c_j) \neq (0, 0, 0)$, $j = 1, 2, 3$; s.t.

$$a||p_i||^2 + b + \sum_{j=1}^3 c_j p_i^j; \quad i = 1, \dots, 5.$$

Then $\Delta_1 = \Delta_2 = 0$, where:

$$\Delta_1(p_1, p_2, p_3, p_4, p_5) = \begin{vmatrix} ||p_1||^2 & 1 & p_1^1 & p_1^2 & p_1^3 \\ ||p_2||^2 & 1 & p_2^1 & p_2^2 & p_2^3 \\ ||p_3||^2 & 1 & p_3^1 & p_3^2 & p_3^3 \\ ||p_4||^2 & 1 & p_4^1 & p_4^2 & p_4^3 \\ ||p_5||^2 & 1 & p_5^1 & p_5^2 & p_5^3 \end{vmatrix}$$

and

$$\Delta_2(p_1, p_2, p_3, p_4, p_5) = \begin{vmatrix} 1 & ||p_1||^2 & -2p_1^1 & -2p_1^2 & -2p_1^3 \\ 1 & ||p_2||^2 & -2p_2^1 & -2p_2^2 & -2p_2^3 \\ 1 & ||p_3||^2 & -2p_3^1 & -2p_3^2 & -2p_3^3 \\ 1 & ||p_4||^2 & -2p_4^1 & -2p_4^2 & -2p_4^3 \\ 1 & ||p_5||^2 & -2p_5^1 & -2p_5^2 & -2p_5^3 \end{vmatrix}$$

Therefore $\Delta_1 \cdot \Delta_2^t = 0$. But $\Delta_1 \cdot \Delta_2^t = \Delta(p_1, p_2, p_3, p_4, p_5)$, so this implication is proven.

³⁰that can be readily generalized to higher dimensions

($\Leftarrow$) $\Delta(p_1, p_2, p_3, p_4, p_5) = 0 \implies \Delta_1 = 0$ and there exist numbers $(a, b, c_j) \neq (0, 0, 0)$, $j = 1, 2, 3$; s.t.

$$a||p_i||^2 + b + \sum_{j=1}^3 c_j p_i^j = 0; \quad i = 1, \dots, 5;$$

i.e. p_1, p_2, p_3, p_4, p_5 belong to the plane or the sphere given by the equation

$$a||X||^2 + b + c \sum_{j=1}^3 c_j X = 0; \quad X = (x_1, x_2, x_3).$$

□

REFERENCES

- [Ba1] Banchoff, T.A. – *Critical points and curvature for embedded polyhedra*, J. Differential Geometry, 1 (1967), 257-268.
- [Ba2] Banchoff, T.A. – *Critical Points and Curvature for Embedded Polyhedral Surfaces*, Amer. Math. Monthly, 77 (1970), 475-485.
- [Be] Berger, M. – *Geometry I*, Universitext, Springer-Verlag, 1987.
- [B] Blumenthal, L. M. *Distance Geometry – Theory and Applications*, Claredon, 1953.
- [BM] Blumenthal, L. M. and Menger, K. – *Studies in Geometry*, Freeman and Co., 1970.
- [BH] Bridson, M. R. and Haefliger, A. – *Metric spaces of non-positive curvature*, Grundlehren der mathematischen Wissenschaften, Springer-Verlag, Berlin, 1999.
- [BBI] Burago, D., Burago, Y. and Ivanov, S. – *A Course in Metric Geometry*, GSM, AMS, RI, 2000.
- [BCM] Borrelli, V. Cazals, F. and Morvan, J.-M. – *On the angular defect of triangulations and the pointwise approximation of Curvatures*, Computer Aided Geometric Designs, 20, pp. 319-341, 2003.
- [CMS] Cheeger, J., Müller, W., and Schrader, R. – *On the Curvature of Piecewise Flat Spaces*, Comm. Math. Phys., 92, 1984, 405-454.
- [C-SM] Cohen-Steiner, D. and Morvan, J.-M. – *Restricted Delaunay triangulations and normal cycle*, preprint, 2003.
- [F] Fu, J. H. G. – *Convergence of Curvatures in Secant Approximation*, J. Differential Geometry, 37, 1993, 177-190.
- [G+] Mikhail Gromov – *Metric structures for Riemannian and non-Riemannian spaces*, Progress in Mathematics 152, Birkhauser, Boston, 1999.
- [K] Kirk, W. A. – *On Curvature of a Metric Space at a Point*, Pacific J. Math. 14: 195-198, 1964.
- [LWZL] Liu, G.H., Wong, Y.S., Zhang, Y.F. and Loh, H.T. – *Adaptive fairing of digitized point data with discrete curvature*, Computer Aided Design, vol. 34(4), 309-320, 2002.
- [MD] Maltret, J.-L. and Daniel, M. – *Discrete curvatures and applications: a survey*, preprint, 2003.
- [P] Pajot, H. – *Analytic Capacity, Rectifiability, Menger Curvature and the Cauchy Integral*, LNM 1799, Springer, Berlin, 2002.
- [RR] Ramsay, A. and Richtmayer, R.D. – *Introduction to Hyperbolic Geometry*, Universitext, Springer-Verlag, 1991.
- [Rat] Ratcliffe, J.C. : *Foundations of Hyperbolic Manifolds*, GTM 194, Springer Verlag, N.Y., 1994.
- [R] Robinson, C.V. – *A Simple Way of Computing the Gauss Curvature of a Surface*, Reports of a Mathematical Colloquium, Second Series, Issue 5-6, 16-24, 1944.
- [SMSER] Surazhsky, T., Magid, E., Soldea, O., Elber, G. and Rivlin, E. – *A Comparison of Gaussian and Mean Curvatures Estimation Methods on TRIangular Meshes*, preprint, 2003.
- [T] Troyanov, M. – *Tangent Spaces to Metric Spaces: Overview and Motivations*, preprint, 2003.

DEPARTMENT OF MATHEMATICS, TECHNION & DEPARTMENT OF SOFTWARE ENGINEERING, ORT
BRAUDE COLLEGE, KARMIEL

E-mail address: `semil@tx.technion.ac.il`